

Centrality of Community Activities in a Network Analysis of Outcome- limiting Factors: Implications for Recovery-oriented Strategies

Armida Mucci

Department of Psychiatry
University of Campania Luigi Vanvitelli

Cognition and Brain imaging Research Group

Department of Psychiatry - University of Campania Luigi Vanvitelli

Silvana Galderisi
Armida Mucci
Paola Bucci
Giuseppe Piegari
Eleonora Merlotti
Annarita Vignapiano
Davide Palumbo
Giulia Maria Giordano
Antonella Amodio
Daria Pietrafesa
Carmen Aiello
Debora Molle
Sara Patriarca
Francesco Brando
Alessia Nicita
Luigi Giuliani
Cristina Romaldi



Campania
University
Luigi Vanvitelli

Armida Mucci Department of Psychiatry

Wien 21-02-2019


Outline of the presentation

- ✓ Implications of recovery-oriented approaches
- ✓ Variables influencing real-life functioning
- ✓ Modeling relationships between disease-related factors and functioning
- ✓ The Italian study on factors influencing real-life functioning in schizophrenia
- ✓ The network analysis
- ✓ Follow-up preliminary results
- ✓ Conclusions

Implications of recovery-oriented approaches


The focus of recovery-oriented approaches is on fostering hope and resilience, fighting self-stigma, supporting self-determination and promoting social inclusion.

The implementation of recovery-oriented plans requires an in depth understanding of key factors influencing real-life functioning.




¹Addington et al, Can J Psychiatry 2003; ¹Harvey & Strassnig World Psychiatry 2013; ¹Galderisi et al, Schizophr Res 2013; ^{1,2,3}Galderisi et al, World Psychiatry; ²Dickerson F, et al. Schizophr Res 1999; ²Green MF, et al. Schizophr Bull 2000; ³Couture et al. Schizophr Bull 2006; ³Green et al, Arch Gen Psychiatry 2012; ³Harvey & Strassnig World Psychiatry 2013; ³Reickmann et al, Schizophr Res 2005.

Negative Symptoms, Cognitive Impairment and Real-Life Functioning



Bowie et al, Am J Psychiatry, 2006

Prediction of participation in everyday life activities



Italian Network for Research on Psychoses (NIRP)

(Coordinator Prof. M. Maj)

- 1 - Sezione di Psichiatria, Dipartimento di Neuroscienze, Università di Torino (*Prof. ssa Rocca*)
- 2 - Struttura Complessa di Psichiatria, Università di Novara (*Prof.ssa Zeppegno*)
- 3 - Cattedra di Psichiatria, Ospedale Maggiore Policlinico, Università di Milano (*Prof. Altamura*)
- 5 - Clinica Psichiatrica, Ospedali Civili, Università di Brescia (*Prof. Vita*)
- 6 - Sezione di Psichiatria, Ospedale S. Martino, Università di Genova (*Prof. Amore*)
- 7 - Sezione di Psichiatria, Dipartimento di Neuroscienze, Università di Padova (*Prof. Santonastaso*)
- 8 - Istituto di Psichiatria, Università di Bologna (*Prof.ssa De Ronchi*)
- 9 - Sezione di Psichiatria, Dipartimento di Neuroscienze, Università di Parma (*Prof. Marchesi*)
- 10 - Clinica Psichiatrica, Università di Firenze (*Prof. Pallanti*)
- 11 - Clinica Psichiatrica, Università di Firenze (*Prof. Ricca*)
- 12 - Clinica Psichiatrica, Presidio Ospedaliero, Università di Pisa (*Prof.ssa Dell'Osso*)
- 13 - Sezione di Psichiatria, Dipartimento di Neuroscienze, Università di Siena (*Prof. Fagiolini*)
- 14 - Clinica Psichiatrica, Ospedale S. Salvatore, Università de L'Aquila (*Prof. ssa Roncone*)
- 15 - Cattedra di Psichiatria, Dipartimento di Medicina Sperimentale, Università de L'Aquila (*Prof. Rossi*)
- 16 - Cattedra di Psichiatria e Centro di Salute Mentale, Università di Chieti (*Prof. di Giannantonio*)
- 17 - Dipartimento di Scienze Psichiatriche e Medicina Psicologica, Università di Roma La Sapienza (*Prof. Biondi*)
- 18 - Cattedra di Psichiatria, II Facoltà di Medicina e Chirurgia, Università di Roma La Sapienza (*Prof. Girardi*)
- 19 - Clinica Psichiatrica, Università di Roma Tor Vergata (*Prof. Siracusano*)
- 20 - Dipartimento di Salute Mentale, Università della Campania L.Vanvitelli (*Prof.ssa Galderisi*)
- 21 - Clinica Psichiatrica, Dipartimento di Neuroscienze, Università Federico II di Napoli (*Prof. De Bartolomeis*)
- 22 - Cattedra di Psichiatria, Università di Salerno (*Prof. Monteleone*)
- 23 - Dipartimento di Scienze Neurologiche e Psichiatriche, Università di Bari (*Prof. Bertolino*)
- 24 - Clinica Psichiatrica, Università di Catania (*Prof. Aguglia*)
- 25 - Cattedra di Psichiatria, Università di Cagliari (*Prof. Carpinello*)
- 26 - Dipartimento di Salute Mentale, Università di Foggia (*Prof. Bellomo*)
- 27 - Unità Operativa Psichiatria Universitaria 2 –Università di Pisa (*Prof. Mauri*)


The influence of illness-related variables, personal resources and context-related factors on real-life functioning of people with schizophrenia

Clinically stable


No relapse or hospitalization

No change in AP treatment

for the three months preceding inclusion




Antipsychotic Treatment



Symptomatic remission 21.8%


Functional remission 12% (32% R; 6% NR)

Final Structural Equation Model



Galderisi et al, World Psychiatry 2014


Relationships between illness-related variables and functioning



Bell et al, *J Nerv Ment Dis.* 2005; Bio & Gattaz, *Schizophr Res.* 2011;
Collip et al, *Schizophr Bull.* 2013; Cella et al, *Schizophr Bull.* 2015

Network analysis

Variable C has least centrality;
A, B and E the highest centrality.



A network analysis of variables influencing real-life functioning

Table 1. Assessment Instruments and Study Measures

| Domains and Variables | Instruments | Measures |
|---|--|--|
| Psychopathologic variables: negative symptoms, depression, overall psychopathologic variables, positive symptoms, disorganization | BNSS ^{17,18} ; CDSS ¹⁹ ; PANSS ²⁰ | BNSS expressive deficit (sum of the subscales blunted affect and alogia), BNSS avolition (sum of the subscales anhedonia, asociality and avolition); CDSS total score; PANSS total, PANSS positive symptom factor (sum of the items delusions, hallucinations, grandiosity and unusual thought content), PANSS disorganization (P2 item; ie, conceptual disorganization) |
| Neurocognition | MCCB ^{21,22} | MCCB domain scores: speed of processing, verbal and spatial learning, reasoning and problem solving, attention, working memory |
| Social cognition | MCCB ^{21,22} ; FEIT ²³ ; TASIT ²⁴ | MSCEIT managing emotion section score; FEIT total; TASIT-1, TASIT-2, and TASIT-3 ^a |
| Real-life functioning | SLOF ²⁵ | SLOF interpersonal relationships, SLOF everyday life skills, SLOF work skills |
| Functional capacity | UPSA-B ²⁶ | UPSA-B total score |
| Service engagement with mental health services | SES ²⁷ | SES total score |
| Internalized stigma | ISMI ²⁸ | ISMI total score |
| Resilience | RSA ²⁹ | RSA factors: perception of self, perception of the future, social competence, family cohesion |
| Incentives | A count variable was created to reflect the availability of a disability pension, access to family financial and practical support, and registration on employment lists | Total incentives |

Abbreviations: BNSS, Brief Negative Symptom Scale; CDSS, Calgary Depression Scale for Schizophrenia; FEIT, Facial Emotion Identification Test; ISMI, Internalized Stigma of Mental Illness; MCCB, Measurement and Treatment Research to Improve Cognition in Schizophrenia Consensus Cognitive Battery; MSCEIT, Mayer-Salovey-Caruso Emotional Intelligence Test; PANSS, Positive and Negative Syndrome Scale; RSA, Resilience Scale for Adults; SES, Service


Engagement Scale; SLOF, Specific Level of Functioning Scale; TASIT, The Awareness of Social Inference Test; UPSA-B, University of California San Diego Performance-based Skills Assessment Brief.

^a TASIT-1 indicates section 1: emotion evaluation; TASIT-2, section 2: social inference minimal; and TASIT-3, section 3: social inference-enriched.

Demographic and clinical characteristics of the study sample


| Characteristic | Total Participants (N = 740) |
|---|------------------------------|
| Age, mean (SD), y | 40.0 (10.9) |
| Male sex, No. (%) | 519 (70.1) |
| Married, No. (%) | 58 (7.8) |
| Working, No. (%) | 216/713 (30.3) |
| Educational level, mean (SD), y | 11.7 (3.3) |
| Age at onset, mean (SD), y | 24.1 (7.1) |
| Duration of illness, mean (SD), y | 16.4 (10.7) |
| Antipsychotic treatment at first episode, No. (%) | |
| First generation | 261 (35.3) |
| Second generation | 236 (31.9) |
| Both | 33 (4.5) |
| Unknown | 210 (28.4) |
| Current antipsychotic treatment (last 3 mo), No. (%) | |
| First generation | 107 (14.5) |
| Second generation | 510 (68.9) |
| Both | 100 (13.5) |
| None | 23 (3.1) |
| Integrated treatment, No. (%) | 203 (27.4) |
| Patients in symptomatic remission, No. (%) ^a | 163 (22.0) |

A network analysis of variables influencing real-life functioning



Resilience: Pse, Perception of self; PFu, Perception of future; SCo, Social competence; FCo, Family cohesion; **Neurocognition:** VLe, Verbal learning; SLe, Visuospatial learning; Att, Attention; PrS, Problem solving; PSp, Processing speed; WMe, Working memory; **Social Cognition:** Ta1-Ta3, TASIT Section 1-3; FEI, FEIT; MSC, MSCEIT managing emotion section; **Real-life Functioning:** Int, SLOF Interpersonal relationships; ELS, SLOF Everyday life skills; Wrk, SLOF work skills; Stg, Stigma; EnS, Service engagement; FC, Functional capacity; Pos, PANSS Positive factor; Dis, PANSS P2-Disorganization; ExD, BNSS Expressive deficit domain; Avl, BNSS Avolition; Dep, CDSS total score for Depression; Inc, Incentives.

A network analysis of variables influencing real-life functioning



Resilience: Pse, Perception of self; PFu, Perception of future; SCo, Social competence; FCo, Family cohesion; **Neurocognition:** VLe, Verbal learning; SLe, Visuospatial learning; Att, Attention; PrS, Problem solving; PSp, Processing speed; WMe, Working memory; **Social Cognition:** Ta1-Ta3, TASIT Section 1-3; FEI, FEIT; MSC, MSCEIT managing emotion section; **Real-life Functioning:** Int, SLOF Interpersonal relationships; ELS, SLOF Everyday life skills; Wrk, SLOF work skills; Stg, Stigma; EnS, Service engagement; FC, Functional capacity; Pos, PANSS Positive factor; Dis, PANSS P2-Disorganization; ExD, BNSS Expressive deficit domain; Avl, BNSS Avolition; Dep, CDSS total score for Depression; Inc, Incentives.

Characteristics of subjects at 4-year follow-up


| Socio-demographic and clinical data - N = 618 | |
|---|-----------|
| Gender (% males) | 69.1 |
| Age (years, mean±SD) | 45.1±10.5 |
| Education (years, mean±SD) | 11.7±3.4 |
| Married (%) | 7.4 |
| Stable affective relationships (%) | 18.9 |
| Working (%) | 34.4 |
| Legal problems (%) | 1.3 |
| Home care (%) | 8.3 |
| Currently in a residential facility (%) | 10.1 |
| Relapses (%) | 43.5 |
| Number of relapses (median, range) | 2 (1-21) |

Remission & recovery

| | N | % |
|----------------------------|---------|------|
| Baseline | | |
| Symptom remission | 140/618 | 22.7 |
| Recovery* | 83/618 | 13.4 |
| 4-year follow-up | | |
| Symptom remission | 185/618 | 29.9 |
| Recovery* | 113/618 | 18.3 |
| | | |
| Recovered at baseline & FU | 36/618 | 5.8 |

*UPSA-B>78 and symptom remission according to Andreasen's criteria
**subjects with complete data on UPSA-B

A network analysis of variables influencing real-life functioning



Resilience: Pse, Perception of self; PFu, Perception of future; SCo, Social competence; FCo, Family cohesion; **Neurocognition:** VLe, Verbal learning; SLe, Visuospatial learning; Att, Attention; PrS, Problem solving; PSp, Processing speed; WMe, Working memory; **Social Cognition:** Ta1-Ta3, TASIT Section 1-3; FEI, FEIT; MSC, MSCEIT managing emotion section; **Real-life Functioning:** Int, SLOF Interpersonal relationships; ELS, SLOF Everyday life skills; Wrk, SLOF work skills; Stg, **Stigma;** EnS, Service engagement; FC, Functional capacity; Pos, PANSS Positive factor; Dis, PANSS P2-Disorganization; ExD, BNSS Expressive deficit domain; Avl, BNSS Avolition; Dep, CDSS total score for Depression; Inc, Incentives.

Galderisi et al JAMA Psychiatry 2018;75(4):396–404..doi:10.1001/jamapsychiatry.2017.4607

Conclusions

The high centrality of functional capacity and everyday life skills suggests that improving the ability to perform tasks relevant to everyday life is critical for any therapeutic intervention in schizophrenia.

The data-driven analysis supports recovery-oriented approaches with a focus on basic functioning goals more than on symptom control.

Special thanks



Italian Network for Research on Psychoses

All subjects with schizophrenia who agreed to participate in the study and all researchers.

Italian Ministry of Education, the Italian Society of Psychopathology, the Italian Society of Biological Psychiatry, Roche, Lilly, AstraZeneca, Lundbeck, and Bristol-Myers Squibb for their financial support to the study.

Thank you for your attention

armida.mucci@gmail.com



PSYCHIATRY IN TRANSITION

TOWARDS
NEW MODELS,
GOALS &
CHALLENGES

SAVE
THE
DATE



EPA 2019
27th
EUROPEAN
CONGRESS
OF PSYCHIATRY

Warsaw, Poland
6-9 April 2019

www.epa-congress.org
#EPA2019